

JOURNAL OF NUMBER THEORY **1**, 500–501 (1969)

## The Punctured Plane is Isomorphic to the Unit Circle

JAMES R. CLAY

*Department of Mathematics, University of Arizona, Tucson, Arizona 85721**Communicated by H. B. Mann*

Received November 13, 1968

This note shows that the multiplicative group of non-zero complex numbers and the multiplicative group of complex numbers of absolute value 1 are isomorphic as groups.

Let  $K^*$  denote the multiplicative group of non-zero complex numbers, and let  $T$  denote the proper subgroup of  $K^*$  of all complex numbers of modulus 1. We shall see that  $K^*$  and  $T$  are isomorphic as groups. First note that  $K^*$  and  $T$  are divisible abelian groups, i.e. each equation  $x^n = b$  has a solution for every positive integer  $n$  and for every element  $b$  of the group. Theorem 19.1 of [3] or Theorem 9.13 of [4] shows that every divisible group is isomorphic to a direct sum of copies of  $\mathbb{Q}$ , the rationals, and copies of quasicyclic groups  $\sigma(p^\infty)$  for various primes  $p$ . Let  $P$  denote the set of all prime natural numbers; thus we have

$$K^* \cong \sum_a \mathbb{Q} \oplus \sum_{p \in P} \sum_{m_p} \sigma(p^\infty),$$

and

$$T \cong \sum_b \mathbb{Q} \oplus \sum_{p \in P} \sum_{n_p} \sigma(p^\infty).$$

We shall show that the cardinal numbers  $a = b$  and each  $m_p = n_p$ . Certainly each  $m_p \geq n_p \geq 1$ . If some  $m_p > 1$ , then  $\sigma(p^\infty) \oplus \sigma(p^\infty)$  is isomorphic to a subgroup of  $K^*$ . Hence, if  $\sigma(p^k)$  is the cyclic group of order  $p^k$ , for some integer  $k > 0$ , then  $\sigma(p^k) \oplus \sigma(p^k)$  is isomorphic to a subgroup of  $K^*$ . So  $K^*$  has  $p^{2k}$  elements satisfying  $x^{p^k} = 1$ , contrary to the fact that there are exactly  $p^k$  elements in  $K^*$  satisfying this equation. Hence  $m_p = n_p = 1$ .

Now  $a = b = 2^{\aleph_0}$ , the first uncountable cardinal, since otherwise  $K^*$  and  $T$  would not have the correct number of elements. So  $K^*$  and  $T$  are each isomorphic to

$$\sum_{2^{\aleph_0}} \mathbb{Q} \oplus \sum_{p \in P} \sigma(p^\infty).$$

It is also of interest to note that  $K^*$  and  $T$  are topological groups with the usual topologies. But  $T$  is compact and  $K^*$  is not.

For more information on groups isomorphic to proper subgroups see [1] and [2].

#### REFERENCES

1. BEAUMONT, R. A. Groups with isomorphic proper subgroups. *Bull. Amer. Math. Soc.* **51** (1945), 381-387.
2. BEAUMONT, R. A. AND PIERCE, R. S. Partially transitive modules and modules with proper isomorphic submodules. *Trans. Amer. Math. Soc.* **91** (1959), 209-219.
3. FUCHS, L. "Abelian Groups." New York, Pergamon, 1960.
4. ROTMAN, J. J. "The Theory of Groups." Boston, Allyn and Bacon, 1965.